



# Tribunil<sup>®</sup>



*[Signature]*  
03/04/2023

**Selective herbicide for weed control  
in wheat, peas, onions, leeks, garlic,  
potatoes and asparagus**

GROUP

5

HERBICIDE

*Contains 700g/kg methabenzthiazuron in the form of  
a wettable powder*



**AGRISOURCE**  
PRODUCTS FOR AGRICULTURE

## WARNING

Keep out of the reach  
of children

Net Contents

**2kg**

**FREE RECYCLING**



**AGRECOVERY**

[www.agrecovery.co.nz](http://www.agrecovery.co.nz)

# READ LABEL COMPLETELY BEFORE USE

## WARNING KEEP OUT OF REACH OF CHILDREN



### HAZARDS

Causes serious eye irritation.

### ECOTOXIC

Very toxic to aquatic life. Very toxic to the soil environment. Do not apply onto or into water. Collect spillage.

## QUALIFICATIONS REQUIRED FOR USE OF THIS CLASS 9 PRODUCT

Anyone who mixes, loads, or applies this product, either by air or ground based methods, is required to be suitably qualified. Refer to Schedule 10 of the Hazardous Substances (Hazardous Property Controls) notice 2017 for a list of suitable qualifications.

## RECORD KEEPING

Where 3 kg or more of the substance is applied within 24 hours, in a place where the substance is likely to enter air or water and leave the application area, a PCBU with management or control of the substance must ensure that a written record is kept of each application of the substance. Refer to Part 4 Clause 48 of the Hazardous Substances (Hazardous Property Controls) notice 2017 for details of information that must be recorded.

## PRECAUTIONS

**Personal Protection:** Do not eat, drink or smoke while using. Wear protective clothing such as impervious gloves, waterproof hat, coat and trousers when using. Remove protective clothing and wash hands and face thoroughly before meals and after work. May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with the eyes.

**Storage:** Store in original container tightly closed and in a locked dry, cool, well ventilated area, away from food stuffs. When stored appropriately this product should show no significant degradation.

**Spills and Disposal:** When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section. Sweep up spilled material, collect and store in properly labelled, sealed drums for safe disposal. Avoid creating dust. If spray mixture is spilled, contain and absorb in suitable inert absorbent material and collect in drums as above. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

**Disposal:** Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container.

**Container Disposal:** Ensure container is completely empty. Recycle through the Agrecovery programme. If this is not possible bury in landfill.

## FIRST AID

If swallowed: Rinse mouth. Do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin contact occurs, remove contaminated clothing and flush skin with running water. If splashed in eyes, wash out immediately with water, if eye or skin irritation persists get medical advice. If inhaled move the victim to fresh air immediately.

**24 HOUR EMERGENCY RESPONSE 0800 CHEMCALL (0800 243 622)**

## DIRECTIONS FOR USE

**Tribunil** is a selective herbicide for the control of various species of weeds in wheat, peas, onions, leeks, garlic, potatoes and asparagus. **Tribunil** is absorbed by both the roots and the leaves of the weeds. As **Tribunil** acts also through the soil a single application gives residual control as well as a knockdown effect. Damage may occur when pre emergence applications are made to light soils, especially under wet conditions.

**NOTE:** Applicators are advised to make sure adequate agitation is used to keep **Tribunil** in suspension in the spray vat.

## WEEDS CONTROLLED

**Susceptible:** Black nightshade, charlock, chickweed, fathen, field speedwell, fumitory, galinsoga, groundsel, henbit, nettle, plantains, annual poa, redroot, scentless chamomile, shepherd's purse, spurrey, small-flowered buttercup, stinking mayweed, twin cress, white campion, wild radish, wild turnip, willow weed.

**Moderately Susceptible:** Combind, redroot (over 4 leaf), sow thistle, wireweed, seedling docks and seedling Californian thistle.

**Resistant:** Cleavers, docks, giant buttercup, rayless chamomile and most deep rooting weeds

## COMPATIBILITY

Do not combine with emulsifiable concentrates of insecticides or other herbicides. Do not use surfactants.

## RESISTANCE MANAGEMENT

Excessive use of any herbicide may lead to the development of resistant weeds. Follow label directions and use in programmes with herbicides from a different action group.

## SPRAY DRIFT

Take all reasonable steps to ensure that **Tribunil** does not cause any significant adverse effects to the environment beyond the application area.

**Wheat: Post Emergence:** Apply 1.0 - 1.5 kg per hectare **Tribunil** + 3.0 litres per hectare of MCPA in 340-350 litres of water from the 5 leaf stage to the shot blade stage. Application should be made as early as possible, however where Californian thistle is a problem application should be delayed until maximum emergence has occurred. Do not apply to wheat crops undersown with legumes.

**Peas: Post Emergence:** Apply 1.5 - 2.0 kg per hectare in 340-600 litres of water preferably when the weeds are at the cotyledon to 2 true leaf stage and the crop is no more than 10cm high. Applications may be made up to the 4 true leaf stage when the crop will be 15cm high. **Tribunil** should not be applied beyond this stage. **Tribunil** 1.0 kg per hectare may be used in conjunction with Sencor® 0.25 - 0.50 kg for control of grass weeds and broadleaf weeds. Use as above.

**Onions: Post Emergence:** Apply 1.5 - 2.0 kg per hectare, or alternatively, use multiple dose technique (2 - 3 sprays of 0.5 - 1.0 kg per hectare 7-10 days apart depending on conditions) in 340 - 450 litres of water. Delay application until the 3 true leaf and apply a directed spray avoiding contact with the bulk of the foliage.

**Leeks: Post Emergence:** Apply 2.0 kg per hectare once plants become established after transplanting, in 340 - 450 litres of water.

**Garlic: Post planting:** Apply 3.0 kg per hectare post planting but before the crop emerges. Post Emergence: Apply 2.0 kg per hectare once the crop has 3 or more true leaves. Multiple dose technique (refer onions) may be used to advantage. Apply 340 - 350 litres of water per hectare.

**Potatoes: Pre-emergence:** Apply 2.5 - 3.0 kg per hectare in 340 - 450 litres of water before the potatoes emerge.

**Asparagus: (Seed beds and 1st year direct seeded):** Apply 2.0 kg per hectare in 340 - 450 litres of water prior to emergence or post emergence when the crop is 10 - 15 cm high. Do not use during the cutting season.

## CROP ROTATION

Residues resulting from the use of **Tribunil** may adversely affect certain susceptible crops. Do not plant crops other than those recommended for treatment on this label within 3 months of applying **Tribunil** without obtaining advice from the distributor. To minimise residue effects, areas treated with **Tribunil** should be ploughed and thoroughly cultivated before sowing following crops.

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

## EXCLUSION OF LIABILITY: PLEASE READ

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Agrisource Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

© Tribunil is a registered trademark of STEFES GmbH

BATCH NO:

Registered pursuant to the ACVM Act 1997, No P1343  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

DATE OF  
MANUFACTURE:

Approved pursuant to HSN0 Act 1996 No HSR000536  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions

## REGISTRANT

STEFES GmbH  
Wendenstr.  
21 b - D - 20097  
Hamburg, Germany

## NZ AGENT

Agri Solutions Ltd  
Level 1, 26 Crummer Road  
Grey Lynn, Auckland 1021

## DISTRIBUTOR

Agrisource (2000) Ltd  
45 Kitchener Road, Pukekohe 2120  
09 237 0442 · [www.agrisource.co.nz](http://www.agrisource.co.nz)



03/04/2023