



FUNGICIDE

# Condor<sup>®</sup>

**For the control of a wide range of leaf and stem diseases of Wheat, Barley, Oats, Ryegrass seed crops and Turf grass and for the treatment of Prairie grass seed for Head Smut**

Contains 250g/L propiconazole in the form of an emulsifiable concentrate

**DANGER**

Keep out of the reach of children

Net Contents **5 LITRES**

GROUP

3

FUNGICIDE



[agrisource.co.nz](http://agrisource.co.nz)



## GENERAL INFORMATION

**Condor** is a systemic fungicide for the control of various foliar diseases of small grain cereals and ryegrass, as well as the treatment of prairie grass seed for Head Smut.

**Wheat:** Stripe Rust, Leaf Rust (Brown Rust), Speckled Leaf Blotch and Glume Blotch

**Barley:** Net Blotch (both net and spot forms), Spot Blotch, Leaf Scald, Leaf Rust and Ovarialia Leaf Spot **Oats:** Crown Rust **Ryegrass:** Stem Rust and Leaf Rust (Crown Rust) **Turf:** Dollar Spot, Brown Patch, Red Thread, Fusarium Patch **Prairie grass (seed treatment):** Head Smut

### RESISTANCE

Resistance to this fungicide and related products could develop from repeated use. To minimise this risk use strictly in accordance with the label instructions. In cereals use a maximum of three DMI applications per crop, including seed treatment, at full recommended label rates.

### APPLICATION

**Maximum application as a seed treatment:** **Condor** must not be applied at rate exceeding 15 g propiconazole per 100 kg seed. **Maximum application rate for wide dissepive use:** **Condor** must not be applied at rates exceeding 125 g propiconazole/ha. **Condor** must not be applied to the same area more than three times in any 365 day period.

**Aerial Application:** Apply in sufficient water to ensure more than 20 droplets per square centimetre on all leaves. As there are many factors involved in aerial application, for optimum results, contact your local supplier representative for detailed information.

### MIXING

Add the required amount of **Condor** to the partly filled spray tank and add the remainder of the water with agitation.

### TREATED SEED

Containers of treated seed must be labelled accordingly. When used as a seed treatment, **Condor** must be applied in a purpose-built seed coating machine, in a treatment plant with local exhaust ventilation. At the time of treatment, seed must be dyed so as to differentiate it from untreated seed. Treated seed must be completely covered by soil when sown. Excess treated seed must not be left in areas accessible to birds. Personal protection equipment must be used when handling seeds treated with **Condor**. Seeds treated with **Condor** must be disposed of in accordance with the disposal requirements for toxic and ecotoxic substances. Seed treated with **Condor** must not be used for human or animal consumption.

### COMPATIBILITY

**Condor** is compatible with most of the commonly used pesticides, including herbicides, used on cereals.

### STORAGE

Store locked up in original container, tightly closed, away from foodstuffs. Store in a cool dry well-ventilated place, avoiding temperatures below minus 5°C or above 35°C.

**In case of fire:** use fine-water spray or foam.

During a fire this product may release smoke and hazardous decomposition products. When fighting a major fire wear an air-supplied respirator.

### DIRECTIONS FOR USE

**It is an offence to use this product on animals.**

### WITHHOLDING PERIODS

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

Cereals (silage and grain crops) ..... 42 days

Ryegrass, Prairie grass seed crops ..... 30 days

Turf ..... DO NOT graze or feed clippings to livestock

### CONDITIONS OF SUPPLY

All goods supplied by us are produced under rigid quality control procedures which assure the purchaser that at the time of sale they are physically suitable for use in the manner stated. However, to the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Agri Solutions Ltd as registrant, nor any other distributor has any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

### REGISTRANT

Agri Solutions Ltd  
45 Kitchener Road,  
Pukekohe 2120

### DISTRIBUTOR

Agrisource (2000) Ltd  
45 Kitchener Road, Pukekohe 2120  
09 237 0442 - www.agrisource.co.nz

### BATCH NO:

### DATE OF

MANUFACTURE:

CROP	DISEASE	RATES	REMARKS
Wheat	Stripe Rust, Speckled Leaf Blotch	500ml in a minimum of 200 litres water/ha	Apply as a preventative spray on susceptible cultivars. For control on less susceptible cultivars, apply at first sign of the diseases. Note: DO NOT apply <b>Condor</b> to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
	Glume Blotch		Apply as a preventative spray at ear emergence on susceptible cultivars especially during cool, moist conditions which favour spread of this disease to the head. On less susceptible cultivars, apply at first sign of the disease. Note: DO NOT apply <b>Condor</b> to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Wheat, Barley	Leaf Rusts	500ml in a minimum of 200 litres water/ha	Apply as a preventative spray at ear emergence on susceptible cultivars. For control on less susceptible cultivars, apply at first sign of the disease. Note: DO NOT apply <b>Condor</b> to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Barley	Net Blotch, Spot Blotch, Leaf Scald, Ovarialia Leaf Spot		Apply at first sign of the diseases. Note: DO NOT apply <b>Condor</b> to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Oats	Crown Rust		Apply as a preventative spray at flag leaf emergence on susceptible cultivars. On less susceptible cultivars, apply at first sign of the disease up to the completion of flowering. Note: DO NOT apply <b>Condor</b> to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Ryegrass seed crops	Stem and Leaf Rusts (Crown Rust)		For optimum control on susceptible cultivars, apply <b>Condor</b> as a preventative spray after closing up for seed, and repeat at 3 to 4 week intervals depending upon continuing disease pressure. For control on less susceptible cultivars, apply at first sign of disease.
Turf Grass	Apply in a minimum of 10 litres of water/100 square metres. Apply as a preventative spray before the signs of disease appear, when conditions are favourable for disease development. For maximum control of turf diseases, maintain the appropriate water and fertiliser requirements to minimise disease development. DO NOT mow within 24 hours of application, and delay application after the morning dew has dried, to minimise the spread of diseases. Note: Some reduction in grass growth may occur. DO NOT apply to newly sown turf.		
	Dollar Spot	20ml/100 square metres	Apply every 14 to 28 days.
	Brown Patch	30ml/100 square metres	Apply every 10 to 21 days. Under conditions of high temperature and humidity, repeat application at the shorter interval.
	Red Thread		Apply every 14 to 21 days.
	Fusarium Patch		Make the first application in late autumn and repeat as necessary, ensuring no more than 3 applications per season.
Prairie grass seed	Head Smut	60ml/100 kg of dressed seed	Apply as a seed coat in sufficient water to ensure adequate coverage. At the time of treatment, the seed must be dyed a distinctive colour (see under 'Treated Seed').

© Condor is a registered trademark of Agri Solutions Limited

# READ LABEL COMPLETELY BEFORE USE

## RECORD KEEPING

Where 3 kg or more of the substance is applied within 24 hours, in a place where the substance is likely to enter air or water and leave the application area, a PCBU with management or control of the substance must ensure that a written record is kept of each application of the substance. Refer to Part 4 Clause 48 of the Hazardous Substances (Hazardous Property Controls) notice 2017 for details of information that must be recorded.

## PCBU RESPONSIBILITIES

A PCBU with management and control of the place to where this product is being used, handled, manufactured, stored or applied must ensure that every worker who handles it, is provided with adequate information, training and instruction. This includes provision and training in the use of any personal protective equipment to be used.

## PERSONNEL QUALIFICATION & EXPERIENCE

This product must be under the control of an appropriate trained and experienced applicator during use. Refer to Schedule 10 of the Hazardous Substances (Hazardous Property Controls) notice 2017 for a list of suitable qualifications.



## HAZARD CLASSIFICATION

Flam. Liquid 4, Eye Irritant 2, Skin Sens. 1, Reproductive toxicity 1, Aquatic Acute 1, Aquatic Chronic 1, Hazardous to soil organisms, Hazardous to terrestrial vertebrates

## WARNING

Combustible liquid. Causes skin irritation. Causes serious eye irritation. May cause an allergic skin reaction. May damage fertility or the unborn child. Very toxic to aquatic organisms with long lasting effects. Hazardous to soil organisms. Hazardous to terrestrial vertebrates.

## HANDLING PRECAUTIONS

Do not handle until all safety precautions have been read and understood. Keep away from flames and hot surfaces. No Smoking. When mixing or applying wear protective clothing, including face shield, gloves and footwear. Chemical resistant PVC-coated gloves must be worn when handling this product. Avoid contact with skin and eyes, and avoid breathing, vapours or spray. If clothing becomes contaminated with product, remove clothing immediately. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse. DO NOT eat, drink or smoke while using. Wash hands and face after handling and before eating, drinking or smoking and after work. Wash protective clothing daily after work.

Apply with well-maintained and calibrated spray equipment. DO NOT re-enter site until spray has dried.

Records of use must be kept. Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.

CONDOR must not be applied into or onto water. Avoid contamination of any water supply with chemical or empty container. Avoid release to the environment. Collect spillage.

**Spillage:** Wear suitable protective clothing including face shield, impervious gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

**Product Disposal:** Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

**Container Disposal:** Ensure the container is empty. Triple rinse the container with water and add the residue to the spray tank. Crush and bury the empty container in an approved local authority facility. DO NOT re-use this container for any other purpose.

**Safety Data Sheet:** Additional information is provided in the SDS available from your supplier.

## FIRST AID

If swallowed DO NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. Begin artificial respiration if the victim is not breathing. Use mouth to nose rather than mouth to mouth. Obtain medical attention. IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. IF INHALED: Move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped. IF exposed or concerned: Get medical advice/attention. If medical advice is needed, have product container or label at hand.

## EMERGENCY RESPONSE

In a transport emergency call 111, Police or Fire Brigade

Registered pursuant to the ACVM Act 1997 No. P9817  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

Approved pursuant to HSNO Act 1996 No. HSR101041  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions

## 24 HOUR EMERGENCY RESPONSE 0800 CHEMCALL (0800 243 622)