



FUNGICIDE

Condor[®]

For the control of a wide range of leaf and stem diseases of Wheat, Barley, Oats, Ryegrass seed crops and Turf grass and for the treatment of Prairie grass seed for Head Smut

Contains 250 g/L propiconazole in the form of an emulsifiable concentrate

DANGER // COMBUSTIBLE LIQUID // ECOTOXIC

Keep out of the reach of children

Net Contents **5 L**

GROUP

3

FUNGICIDE



agrisource.co.nz



GENERAL INFORMATION

Condor is a systemic fungicide for the control of various foliar diseases of small grain cereals and ryegrass, as well as the treatment of prairie grass seed for Head Smut.

IMPORTANT - RESISTANCE MANAGEMENT

Condor is a **GROUP 3 FUNGICIDE** from the Imidazole (DMI/Triazole) group of fungicides. Resistance to Condor and related DMI fungicides could develop from repeated use. To minimise this risk, use strictly in accordance with label instructions. Apply no more than 3 DMI treatments per season including seed treatment.

It is an offence to use this product on animals.

APPLICATION

Maximum application as a seed treatment: Condor must not be applied at rate exceeding 15 g propiconazole per 100 kg seed.

Maximum application rate for wide dispersive use: Condor must not be applied at rates exceeding 125 g/ha of propiconazole. Condor must not be applied to the same area more than three times in any 365 day period.

Aerial Application: Apply in sufficient water to ensure more than 20 droplets per square centimetre on all leaves. As there are many factors involved in aerial application, for optimum results, contact your local supplier representative for detailed information.

MIXING

Add the required amount of Condor to the partly filled spray tank and add the remainder of the water with agitation.

TREATED SEED

Containers of treated seed must be labelled accordingly. When used as a seed treatment, Condor must be applied in a purpose-built seed coating machine, in a treatment plant with local exhaust ventilation. At the time of treatment, seed must be dyed so as to differentiate it from untreated seed. Treated seed must be completely covered by soil when sown. Excess treated seed must not be left in areas accessible to birds. Personal protection equipment must be used when handling seeds treated with Condor.

For treated seeds, information must be provided that details the substance that the seeds have been treated with, its hazardous properties, and precautions to be taken in handling the treated seeds.

Disposal: Seeds treated with Condor shall be disposed of in accordance with clauses 9 and 10 of the Hazardous Substance (Disposal) Notice 2017. Seed treated with Condor must not be used for human or animal consumption.

COMPATIBILITY

Condor is compatible with most of the commonly used pesticides, including herbicides, used on cereals.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS

Cereals (silage and grain crops) 42 days
Ryegrass, Prairie grass seed crops 30 days
Turf DO NOT graze or feed clippings to livestock

CONDITIONS OF SUPPLY

All goods supplied by us are produced under rigid quality control procedures which assure the purchaser that at the time of sale they are physically suitable for use in the manner stated. However, to the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Agri Solutions Ltd as registrant, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

*Condor is a registered Trademark of Agri Solutions Ltd

ACVM Registration No. P009817
See www.foodsafety.govt.nz for registration conditions

HSNO Approval No. HSR101041
See www.epa.govt.nz for approval conditions

REGISTRANT
Agri Solutions Ltd
45 Kitchener Road,
Pukekohe 2120

DISTRIBUTOR
Agrisource (2000) Limited
45 Kitchener Road, Pukekohe 2120
08 237 0422 - www.agrisource.co.nz

BATCH NO:

DATE OF
MANUFACTURE:

CROP	RATE	RATES	REMARKS
Wheat	Stripe Rust, Speckled Leaf Blotch		Apply as a preventative spray on susceptible cultivars. For control on less susceptible cultivars, apply at first sign of the diseases. Note: DO NOT apply Condor to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
	Glume Blotch		Apply as a preventative spray at ear emergence on susceptible cultivars especially during cool, moist conditions which favour spread of this disease to the head. On less susceptible cultivars, apply at first sign of the disease. Note: DO NOT apply Condor to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Wheat, Barley	Leaf Rusts	500 mL in a minimum of 200 L water/ha	Apply as a preventative spray at ear emergence on susceptible cultivars. For control on less susceptible cultivars, apply at first sign of the diseases. Note: DO NOT apply Condor to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Barley	Net Blotch, Spot Blotch, Leaf Scald, Ovariala Leaf Spot		Apply at first sign of the diseases. Note: DO NOT apply Condor to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Oats	Crown Rust		Apply as a preventative spray at flag leaf emergence on susceptible cultivars. On less susceptible cultivars, apply at first sign of the disease up to the completion of flowering. Note: DO NOT apply Condor to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Ryegrass seed crops	Stem and Leaf Rusts (Crown Rust)		For optimum control on susceptible cultivars, apply Condor as a preventative spray after closing up for seed, and repeat at 3 to 4 week intervals depending upon continuing disease pressure. For control on less susceptible cultivars, apply at first sign of disease.
Turf Grass			Apply in a minimum of 10 L of water/100 m ² . Apply as a preventative spray before the signs of disease appear, when conditions are favourable for disease development. For maximum control of turf diseases, maintain the appropriate water and fertiliser requirements to minimise disease development. DO NOT mow within 24 hours of application, and delay application to after the morning dew has dried, to minimise the spread of diseases. Note: Some reduction in grass growth may occur. DO NOT apply to newly sown turf.
	Dollar Spot	20 mL/100 m ²	Apply every 14 to 28 days.
	Brown Patch		Apply every 10 to 21 days. Under conditions of high temperature and humidity, repeat application at the shorter interval.
	Red Thread	30 mL/100 m ²	Apply every 14 to 21 days.
	Fusarium Patch		Make the first application in late autumn and repeat as necessary, ensuring no more than 3 applications per season.
Prairie grass seed	Head Smut	60 mL/100 kg of dressed seed	Apply as a seed coat in sufficient water to ensure adequate coverage. At the time of treatment, the seed must be dyed a distinctive colour (see under 'Treated Seed').

READ LABEL COMPLETELY BEFORE USE



DANGER

Causes serious eye irritation. May cause an allergic skin reaction. May damage fertility or the unborn child.

COMBUSTIBLE LIQUID

Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No Smoking.

ECOTOXICITY

Very toxic to aquatic life with long lasting effects. Do not apply onto or into water. Avoid contamination of any water supply with product or empty container. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Collect spillage. Hazardous to terrestrial vertebrates. Take all reasonable steps to ensure non-target terrestrial vertebrates are not adversely affected by this product.

RECORD KEEPING

Where 3 kg or more of the substance is applied within 24 hours, in a place where the substance is likely to enter air or water and leave the application area, a PCBU with management or control of the substance must ensure that a written record is kept of each application of the substance. Refer to Part 4 Clause 48 of the Hazardous Substances (Hazardous Property Controls) notice 2017 for details of information that must be recorded.

PCBU RESPONSIBILITIES

A PCBU with management and control of the place to where this product is being used, handled, manufactured, stored or applied must ensure that every worker who handles it, is provided with adequate information, training and instruction. This includes provision and training in the use of any personal protective equipment to be used.

QUALIFICATIONS REQUIRED FOR APPLICATION OF CLASS 9 SUBSTANCES

Anyone who mixes, loads, or applies this product, either by air or ground based methods, is required to be suitably qualified. Refer to Schedule 10 of the Hazardous Substances (Hazardous Property Controls) notice 2017 for a list of suitable qualifications.

HANDLING PRECAUTIONS - FOR WORKPLACE USE ONLY

Do not handle until all safety precautions have been read and understood. Do not breathe fumes, vapours or spray. Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. Do not eat, drink or smoke when using this product. When mixing or applying, avoid contact with skin and eyes. Wear full protective clothing, gloves and goggles. Wash hands and face before meals and after work. Wash protective clothing after work. Contaminated work clothing should not be allowed out of the workplace.

Equipment: Apply using accurately calibrated and maintained equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409). Flush spray equipment thoroughly with water after use.

Storage: Store in original container tightly closed and in a locked, dry, cool, well-ventilated area away from fertilisers, feed, seeds and foodstuffs, avoiding temperatures below minus 5°C or above 35°C. Keep out of reach of children. Store in accordance with NZ 8409 Management of Agrichemicals. Stores containing more than 100 L of **Condor** are subject to signage and secondary containment and require an emergency response plan.

Shelf Life: When stored appropriately this product should show no significant degradation for two years from the date of manufacture.

Spillage: Extinguish any ignition sources. Move non-essential people from the area. Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Sweep up solids without creating dust. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

Product Disposal: Dispose of unwanted product by use by another operator strictly in accordance with label directions or through an approved facility.

Treated Seed Disposal: Seeds treated with **Condor** shall be disposed of in accordance with clauses 9 and 10 of the Hazardous Substance (Disposal) Notice 2017

Container Disposal: Triple rinse container and add rinsate to spray tank. Dispose of cleaned container at your local AGRECOVERY container collection site.

Safety Data Sheet: Additional information is provided in the SDS available from your supplier or www.agrisource.co.nz

FIRST AID

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing. IF SWALLOWED: Rinse Mouth. DO NOT induce vomiting. Immediately call a POISON CENTER or doctor/physician. For advice, contact the National Poisons Centre 0800 Poison (0800 764 766) or a doctor immediately. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. IF ON SKIN: Take off contaminated clothing and wash before reuse. Wash with plenty of soap and water. Get medical advice/attention if you feel unwell. If medical advice is needed, have product container or label at hand.

EMERGENCY RESPONSE

In case of fire: Use dry chemical, water spray or carbon dioxide for extinction. Contain spillage. In a transport emergency call 111, Police or Fire Brigade.

24 HOUR EMERGENCY RESPONSE 0800 CHEMCALL (0800 243 622)