



HERBICIDE

Tribunil®

**Selective herbicide for weed control in
wheat, peas, onions, leeks, garlic,
potatoes and asparagus**

Contains 700g/kg methabenzthiazuron in the
form of a wettable powder

WARNING // ECOTOXIC

Keep out of the reach of children

Net Contents **2 KG**

GROUP



5

HERBICIDE

agrisource.co.nz



READ LABEL COMPLETELY BEFORE USE

 	<p>WARNING Causes serious eye irritation.</p> <p>ECOTOXIC Very toxic to aquatic life. Very toxic to the soil environment. Do not apply onto or into water. Collect spillage.</p>
---	--

RECORD KEEPING
Where 3 kg or more of the substance is applied within 24 hours, in a place where the substance is likely to enter air or water and leave the application area, a PCBU with management or control of the substance must ensure that a written record is kept of each application of the substance. Refer to Part 4 Clause 48 of the Hazardous Substances (Hazardous Property Controls) notice 2017 for details of information that must be recorded.

QUALIFICATIONS REQUIRED FOR APPLICATION OF SUBSTANCES HAZARDOUS TO THE ENVIRONMENT
Anyone who mixes, loads, or applies this product, either by air or ground based methods, is required to be suitably qualified. Refer to Schedule 10 of the Hazardous Substances (Hazardous Property Controls) notice 2017 for a list of suitable qualifications.

HANDLING PRECAUTIONS - FOR WORKPLACE USE ONLY

Personal Protection: Do not handle until all safety precautions have been read and understood. Do not breathe the mist, or spray. Do not eat, drink or smoke when using this product. When mixing or applying, avoid contact with skin and eyes. Wear full protective clothing, gloves and goggles. Wash hands and face before meals and after work. Wash protective clothing after work. Contaminated work clothing should not be allowed out of the workplace.

Equipment: Apply using accurately calibrated and maintained equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409). Flush spray equipment thoroughly with water after use.

Storage: Store in original container tightly closed and in a locked, dry, cool, well ventilated area away from fertilisers, feed, seeds and foodstuffs. Keep out of reach of children. Store in accordance with NZ 8409 Management of Agrichemicals.

Shelf Life: When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

Spillage: Move non-essential people from the area. Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Sweep up solids without creating dust. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

Product Disposal: Dispose of unwanted product by use by another operator strictly in accordance with label directions or at an approved facility.

Container Disposal: Ensure container is completely empty. Triple rinse container and add rinsate to mix tank. Recycle cleaned container through the Agrecovery programme. If this is not possible bury in landfill.

Safety Data Sheet: Additional information is provided in the SDS available from your supplier or www.agrisource.co.nz

FIRST AID
IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing. IF SWALLOWED: Rinse Mouth. DO NOT induce vomiting. Call a POISON CENTER or doctor/physician if you feel unwell. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. IF ON SKIN: Wash with plenty of soap and water. Take off contaminated clothing and wash before reuse. If skin irritation or rash occurs: Get medical advice/attention. IF exposed or concerned: Get medical advice/attention. If medical advice is needed, have product container or label at hand. For advice, contact the National Poisons Centre 0800 Poison (0800 764 766).

EMERGENCY RESPONSE
In a transport emergency call 111, Police or Fire Brigade.

® Tribunil is a registered trademark of STEFES GmbH

Registered pursuant to the ACVM Act 1997, No P1343
See www.foodsafety.govt.nz for registration conditions

Approved pursuant to the HSNO Act 1996 No HSR000536
See www.epa.govt.nz for approval conditions

REGISTRANT
STEFES GmbH
Wendenstr. 21 A · 20097
Hamburg, Germany

NZ AGENT
Agri Solutions Ltd
45 Kitchener Road, Pukekohe 2120

DISTRIBUTOR
Agrisource (2000) Ltd
45 Kitchener Road, Pukekohe 2120
09 237 0442 · www.agrisource.co.nz

BATCH NO:

DATE OF
MANUFACTURE:

GENERAL INFORMATION

Tribunil is a selective herbicide for the control of various species of weeds in wheat, peas, onions, leeks, garlic, potatoes and asparagus. **Tribunil** is absorbed by both the roots and the leaves of the weeds. As **Tribunil** acts also through the soil a single application gives residual control as well as a knockdown effect. Damage may occur when pre emergence applications are made to light soils, especially under wet conditions.

NOTE: Applicators are advised to make sure adequate agitation is used to keep **Tribunil** in suspension in the spray vat.

IMPORTANT - RESISTANCE MANAGEMENT

Tribunil contains the **GROUP 5 HERBICIDE** methabenzthiazuron with the inhibition of photosynthesis at photosystem II mode of action. Some naturally occurring weed biotypes that are resistant to triazine herbicides are known to exist. These resistant biotypes may dominate when **Tribunil** or other related Group 5 Herbicides are used excessively. To minimise this risk use strictly in accordance with label instructions and resistance management strategies. See www.resistance.nzpps.org for further information.

WEEDS CONTROLLED

Susceptible: Black nightshade, charlock, chickweed, fathen, field speedwell, fumitory, galinsoga, groundsel, henbit, nettle, plantains, annual poa, redroot, scentless chamomile, shepherd's purse, spurrey, small-flowered buttercup, stinking mayweed, twin cress, white campion, wild radish, wild turnip, willow weed.

Moderately Susceptible: Cornbind, redroot (over 4 leaf), sow thistle, wireweed, seedling docks and seedling Californian thistle.

Resistant: Cleavers, docks, giant buttercup, rayless chamomile and most deep rooting weeds

COMPATIBILITY

Do not combine with emulsifiable concentrates of insecticides or other herbicides. Do not use surfactants.

DIRECTIONS FOR USE

SPRAY DRIFT

Take all reasonable steps to ensure that **Tribunil** does not cause any significant adverse effects to the environment beyond the application area.

Crop	Rate	Application
Wheat	1.0-1.5 kg/ha + 3.0 L/ha of MCPA in 340-350L of water/ha	Post Emergence: Apply from the 5 leaf stage to the shot blade stage. Application should be made as early as possible, however where Californian thistle is a problem application should be delayed until maximum emergence has occurred. Do not apply to wheat crops under sown with legumes.
Peas	1.5-2.0 kg/ha in 340-600L of water/ha	Post Emergence: Apply preferably when the weeds are at the cotyledon to 2 true leaf stage and the crop is no more than 10 cm high. Applications may be made up to the 4 true leaf stage when the crop will be 15cm high. Tribunil should not be applied beyond this stage. Tribunil 1.0 kg/ha may be used in conjunction with metribuzin 480 g/L at 0.25-0.50 kg/ha for control of grass weeds and broadleaf weeds. Use as above.
Onions	1.5-2.0 kg/ha in 340-450L of water/ha	Post Emergence: Alternatively, use multiple dose technique (2-3 sprays of 0.5-1.0 kg/ha 7-10 days apart depending on conditions). Delay application until the 3 true leaf and apply a directed spray avoiding contact with the bulk of the foliage.
Leeks	2.0 kg/ha in 340-450L of water/ha	Post Emergence: Apply once plants become established after transplanting.
Garlic	2.0 kg/ha in 340-350L of water/ha	Post Emergence: Apply once the crop has 3 or more true leaves. Multiple dose technique (refer onions) may be used to advantage.
	3.0 kg/ha in 340-350L of water/ha	Post planting: Apply post planting but before the crop emerges.
Potatoes	2.5-3.0 kg/ha in 340-450L of water/ha	Pre-emergence: Apply before the potatoes emerge.
Asparagus	2.0 kg/ha in 340-450L of water/ha	Seed beds and 1st year direct seeded: Apply prior to emergence or post emergence when the crop is 10-15cm high. Do not use during the cutting season.

CROP ROTATION

Residues resulting from the use of **Tribunil** may adversely affect certain susceptible crops. Do not plant crops other than those recommended for treatment on this label within 3 months of applying **Tribunil** without obtaining advice from the distributor. To minimise residue effects, areas treated with **Tribunil** should be ploughed and thoroughly cultivated before sowing following crops.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

EXCLUSION OF LIABILITY: PLEASE READ

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Agrisource Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

24 HOUR EMERGENCY RESPONSE 0800 CHEMCALL (0800 243 622)